

Conventional Plays

Upper Cretaceous-Lower Tertiary Play

(USGS 4101)

General Characteristics

This hypothetical, continuous-type "tight-gas" play is largely restricted to the marginal marine, partly deltaic Trinidad Sandstone. Although, stratigraphic traps could occur in the Vermejo and Raton Formations (Fig. P-37). Dolly and Meissner (1977) estimated the uppermost Cretaceous/lowermost Tertiary section may have generated approximately 23 TCFG, of which approximately 6 TCFG may be recoverable. Additionally, as much as 750 BCFG of recoverable gas reserves exists in basin-centered gas in the northern portion of the Raton Basin.

**Reservoirs:** The Cretaceous Trinidad Sandstone, characterized as marginal marine and partly deltaic, is the potential reservoir rock. General thickness probably varies between 100 and 250 feet. Reservoir sandstones may be as much as 50 feet thick and reservoir porosity is probably 10-14 percent, but porosity usually varies between 2 and 18 percent. Other potential clastic reservoirs include the Cretaceous Vermejo and Cretaceous/Paleocene Raton Formations.

**Source rocks, timing, and migration:** Pierre Shale and coal/carbo-naceous beds are potential sources in the Vermejo, Raton, and Poison Canyon Formations. Generation and migration probably began no earlier than Eocene.




Figure P-37. Upper Cretaceous-Lower Tertiary Play (4101) with respect to the Pueblo Indian Reservations (modified after Gautier et al., 1996).

**Traps:** Trinidad Sandstones, in a basin-center environment, may lack a conventional seal. Depth to production is rather shallow, ranging between approximately 4,000 and 6,000 feet.

**Exploration status and resource potential:** The play is poorly explored. It is possible that a few new discoveries will exceed 6 BCFG. A number of small gas fields possibly could be found.

Jurassic-Lower Cretaceous Play

(USGS 4102)

General Characteristics

This is a high-risk play, and potential reservoirs are restricted to Jurassic Morrison and Cretaceous Dakota Sandstones deposited as highly lenticular marine bars and fluvial channels (Fig. P-38). Sandstones may be fine to coarse grained, 10-40 feet thick and log-derived porosity may reach 15-25 percent. Field pressure, determined from the now-abandoned Wagon Mound Field (Mora County, New Mexico) is low.

**Reservoirs:** Jurassic Morrison and Cretaceous Dakota Sandstones, deposited as highly lenticular marine bars and fluvial channels, are the potential reservoirs. Sandstones are fine to coarse grained, 10-40 feet thick. Porosity, determined from logs (Figs. 39 and 40), varies between 15 and 25 percent and permeability appears high. Field pressure is low (5.5 psi). Most of the gas has been found in the upper Dakota Sands.




Figure P-38. Jurassic-Lower Cretaceous Play (4102) with respect to the Pueblo Indian Reservations (modified after Gautier et al., 1996).

**Source rocks, timing, and migration:** Shale and coal are possible sources within the Purgatoire-Dakota sequence, and overlying shales include potential source beds for oil and gas. Generation probably began in early Tertiary (Eocene-time) when overlying strata were at least 10,000 feet thick. Migration probably began in the Eocene.

**Traps:** Gas was structurally trapped in the Dakota Sands in a low-relief, northeast-trending Laramide Anticline. Some gas was trapped in lenticular, fluvial Jurassic Morrison Sandstones. Interbedded shales probably act as traps. Depth to known occurrences is 500-5,000 feet.

**Exploration status:** The play is poorly explored. It is unlikely that new discoveries will exceed 6 BCFG or 1 MMBO. A number of small gas fields could probably be found.

**Resource potential:** This is a high risk play; undiscovered resources are estimated to be of small size.

Figure P-40. Typical electric log characteristics of the Dakota Sandstone in the Raton Basin, Colfax County, New Mexico (modified after Gilbert and Asquith, 1976).





Figure P-39. Typical gamma-ray log of Dakota Formation in south-central Raton Basin (modified after Gilbert and Asquith, 1976).



## UNCONVENTIONAL PLAYS

### Coal-Bed Gas Plays

Three coal-bed gas plays are identified in the Raton Basin Province. The Southern Raton Basin Play (4152) is relevant to the Pueblo Indian Reservations (Fig. P-41). Tyler et al., 1991; Stevens et al., 1992; and Close and Dutcher, 1993, have described the geologic controls and potential of coal-bed gas in the Raton Basin, southeastern Colorado and northeastern New Mexico.

In the Raton Basin, coal beds with potential for coal-bed gas are contained within the Upper Cretaceous Vermejo and Upper Cretaceous-Paleocene Raton Formations (Fig. P-42). The Vermejo Formation is as much as 350 feet thick and individual coal seams are as much as 14 feet thick. The cumulative coal thickness for the formation ranges from 5 to 35 feet. The overlying Raton Formation is as much as 1,600 feet thick and has a net coal thickness in the range of 10 to 120 feet. Although the Raton Formation contains more coal, individual coal seams are thinner, more discontinuous, and distributed over 1,200 feet of section. The nature of the coal seams in the two formations is controlled by depositional environment; the Vermejo was deposited in a lagoonal environment; whereas, the Raton was deposited in a fluvial setting. Although coal beds are as much as 4,100 feet deep in the northern part of the basin along the LaVeta Syncline, they are generally less than 1,200 feet over a large part of the basin.

The rank of coals in the Vermejo Formation ranges from high-volatile C bituminous along the margins of the basin to low-volatile bituminous in the central part of the basin. The rank generally coincides with present-day depth of burial and structural configuration, and probably resulted from maximum depth of burial that occurred in early Tertiary time. However, the highest ranks (low-volatile bituminous) occur along the eastward-flowing Purgatoire River where the present-day depths of burial are less than about 1,200 feet. These high ranks are interpreted to be the result of high heat flow from the crust, upper mantle, and (or) deep igneous intrusions which was transferred laterally by groundwater flow in middle Tertiary time. During this time, Vermejo and Raton coal beds commonly served as planes of weakness for igneous intrusions. However, the thermal maturity of the coal beds is only locally affected (one-dike width) by the intrusions.

Coal-bed gases from production tests in the Raton Basin are composed mostly of methane with minor amounts of ethane, carbon dioxide, and nitrogen (each less than 1 percent). Isotopic analyses indicate that the gases are predominantly of thermogenic origin and were probably generated during time of maximum burial and (or) heat flow. Some mixing of relatively recent biogenic gas may occur in areas of groundwater flow.


The Raton Basin is a strongly asymmetric basin with a gently dipping eastern flank and a steeply dipping western flank that is thrust-faulted (Fig. P-9). Several major folds are located along the western margin of the basin. Minor normal faulting occurs within the basin with displacements generally less than 50 feet. The primary fracture permeability system in both the coals and adjoining

rocks is oriented east-west.

Groundwater recharges the topographically high Vermejo-Raton Aquifer along the western margin of the basin and flows eastward to discharge at the topographically lower eastern outcrops along major seeps and drainage areas (Geldon, 1989; Stevens et al., 1992). The primary fracture (face cleat) trend in the Raton Basin coals is perpendicular to the local trend of the Sangre de Cristo thrust front and thus enhances groundwater flow and the potential for artesian overpressuring (Tyler et al., 1991).

Gas contents of coal beds in the basin are highly variable and range from 4 to 810 Scf/t. These contents seem to correlate more closely with depth below the hydrologic potentiometric surface than with depth below the ground surface. On the basis of coal thickness, coal density, drillable area, and gas content, in-place coal-bed gas resources of the Raton Basin are estimated to be as much as 12 TCF (Fig. P-42).

Some coal is produced by both underground and surface methods in the Colorado and New Mexico parts of the basin. Mine-related emissions are minor. An explosion in an underground mine near Trinidad, Colorado, indicates that the coal beds are gassy. Since the late 1970's, more than 110 exploration wells have been drilled for coal-bed gas in the Raton Basin, both in Colorado and New Mexico. Production tests have been variable, but gas rates of more than 300 MCF/D have been reported. At present, all wells are shut-in because of the absence of gas pipelines in the basin. However, a pipeline is under construction and a pilot nitrogen injection project for coalbed gas wells has been approved (Rice and Finn, 1995).




**<- Figure P-41.** Structure map of the Sheep Mountain Field showing the outline of the Sheep Mountain unit. Datum is the top of the Dakota Sandstone Formation (modified after Roth, 1983).

**Figure P-42. ->** Raton Basin coal-bed methane in place, contoured in 4 BCF/mi intervals; data inside the 16 BCF/mi contour have a high degree of uncertainty (modified after Stevens et al., 1992).

## Analog Field Within Raton Basin (near Pueblo Indian Reservations)

### Sheep Mountain Field (Figure P-41)

Location of Discovery Well:	No. 1 Faris, nw se 15, T25S, R70W
Producing Formation:	Cretaceous Dakota Ss; Jurassic Entrada Ss
Type of Trap:	Stratigraphic (shale)
Initial Production:	4900 MCFD
Cumulative Production:	NR
Gas Characteristics:	NR
Type of Drive:	Gas pressure
Average Net Pay:	350 feet, 110 feet
Porosity:	NR
Permeability:	NR






# Southern Raton Basin Play

(USGS 4152)

## General Characteristics

The target area for coal-bed gas is where coal beds of the Vermejo and Raton Formations are greater than 500 feet deep. The thicker, more continuous seams of the Vermejo Formation are probably better targets for coal-bed gas production. The Southern Raton Basin Play target area (Fig. P 43) is based on depth, coal rank, and concentration of gas in place. Exploration wells have been drilled for coal-bed gas in the play, but production has not been established (as of 1996). The reserve potential of coal-bed gas from all three plays is considered very good, but production will depend on infrastructure development, particularly pipeline construction.

In the Southern Raton Basin Play (4152), coal ranks are as much as medium-volatile bituminous, but depths of burial are less than 1,400 feet. Because of these relatively shallow depths, concentrations of gas in-place are about 8 BCF/square mile or less. The reserve potential of this play is also regarded as good (Rice and Finn, 1995).



# South-Central New Mexico Province

This frontier petroleum province covers about 39,900 square miles, primarily in the easternmost part of the Basin and Range Physiographic Province; it has no production (Fig. P-44). For a more complete description of this province, see Butler, 1988; and Grant and Foster, 1989.

Small, northeast-trending rift basins are the predominant physiographic feature of the South-Central New Mexico Province. As much as 10,000 feet of alluvium and volcanic rocks fill these extensional basins, obscuring a moderately thick section of Paleozoic strata.

This province has a complex geologic history, having been deformed by three major periods of tectonism during Phanerozoic time: (1) Late Paleozoic formation of the Ancestral Rocky Mountains, (2) Laramide compression, and (3) Cenozoic relaxation and extension and volcanism. South-Central New Mexico was near the terminus of the northeast-trending transcontinental basement arch during the Late Proterozoic and Paleozoic. Within this time span, sediments were deposited in platform, shallow shelf, basinal, and alluvial plain environments. Epeiric seas generally transgressed from the south, and

thus a greater thickness of strata was deposited during this time in the southern part of the province. During the mid-Paleozoic, general quiescence of the craton in the equatorial paleolatitudes resulted in widespread deposition of fossiliferous carbonates accompanied by basin-margin organic buildups. Convergence of the North and South American tectonic plates in the late Paleozoic resulted in intraplate deformation.


Triassic and Jurassic strata are not well represented in the province, which depositio nally represents an erosional surface shifting from highlands to interior lowlands and coastal plains. Nascent opening of the Gulf of Mexico (Chihuahua Trough) deposited as much as 750 feet of marine Jurassic sediments in the southernmost Mesilla Basin. Continued opening near the New Mexico-Mexico border resulted in an east-west Early Cretaceous rift, extending into southeastern Arizona. A thick Late Cretaceous section of marine sandstones and shales and continental fluvial clastics and paludal coals was deposited as seas transgressed and regressed from the north-northeast and from the south-southwest; about 3,000 feet of this sec

**Figure P-43.** Southern Raton Basin Coal-Bed Play (4152), with respect to the Pueblo Indian Reservations (modified after Gautier et al., 1996).

tion is preserved. Laramide compression from the southwest rejuvenated older fault-bounded structures and other paleo-zones of weakness (for example thrust faults) and created basement-cored uplifts. Plutons, with attendant rich mineralization, intruded the province. Early Tertiary uplift provided cyclic alluvial-fluvial fan gravels and deltaic clastics to continental interior-drained basins and small lakes.

Two hypothetical conventional plays were assessed in this

province. They are Orogrande Basin Play (2602) and Mesilla-Mimbres Basins Play (2603), neither of which occur in or near the Pueblo Indian Reservations. However, limited exploration has occurred within the Acoma Basin Play, which underlies a segment of the Pueblo Reservation. A brief description of the production within the Acoma Basin is presented in the following section.



**Figure P-44.** Outline of South-Central New Mexico Geologic Province with exploration wells from 1900-1993 illustrated. The Acoma Basin is highlighted (modified after Gautier et al., 1996).

Acoma Basin Play

The Acoma Basin has seen limited exploration since the 1920's (Fig. P-45). The boundary between the Acoma Basin and the San Juan Basin to the northwest and the eastern Baca Basin is transitional. Three significant exploratory wells have been drilled in the Acoma Basin since 1981. The Topaz Southwest Number 1 State tested Cretaceous sandstones and shales and Jurassic sandstones with no reported shows. The Austra-Tex Numbers 1 through 7 Rio Puerco Federal, were drilled to test the Pennsylvanian section (Fig. P-46). Primary reservoir targets in the Acoma Basin are Permian lime stones and sandstones of the San Andres and Yeso Formations and Pennsylvanian sandstone and limestones (Fig. P-47). The Cretaceous

Mesaverde and Dakota Sandstones are secondary reservoir targets that have been eroded from the eastern part of the basin and are present only at shallow depths (less than 1500 feet {Broadhead, 1989}) in the western part of the basin (Fig. P-48). Pennsylvanian and Cretaceous marine shales are potential source rocks, but the Cretaceous section may be thermally immature (Broadhead, 1989).

Figure P-45. Outline of the South-Central New Mexico Province with the Acoma Basin Play Highlighted (modified after Gautier et al., 1996).





Figure P-48. Location of exploration wells in Acoma Basin (modified after Gautier et al., 1996.)




Figure P-46. Stratigraphic section depicting bedding relationships within the South-Central New Mexico Geologic Province (modified after Gautier et al., 1996).




Figure P-47. Stratigraphic cross-section E-E' of the Acoma Basin (Fig. P-7; cross-section 5) (modified after Molenaar, 1974).



Pueblo Reservations of New Mexico  
REFERENCES

Armstrong, R.L., 1968, The Sevier orogenic belt in Nevada and Utah: Geological Society of America Bulletin, v. 79, p. 229-258.

Baltz, E.H., 1965, Stratigraphy and history of Raton Basin and notes on San Luis Basin, Colorado-New Mexico: AAPG Bulletin, v. 49, p. 2041-2075.

Baltz, E.H., 1967, Stratigraphy and regional tectonic implications of part of Upper Cretaceous and Tertiary rocks, east-central San Juan Basin, New Mexico: U.S. Geological Survey Professional Paper 552, 101p.

Bachman, G.O., 1953, Geology of a part of northwestern Mora County, New Mexico: U.S. Geological Survey, Oil and Gas Investigations Map OM-137.

Black, B.A., and Hiss, W.L., 1974, Structure and stratigraphy in the vicinity of the Shell Oil Co., Santa Fe Pacific No.1 test well, southern Sandoval County, New Mexico, in Siemers, C.T., ed., Ghost Ranch, central-northern New Mexico: New Mexico Geological Society 25th Field Conference Guidebook, p. 365-370.

Black, B.A., 1982, Oil and gas exploration in the Albuquerque Basin, in Grambling, J.A., and Wells, S.G., eds., Albuquerque country II: New Mexico Geological Society 33rd Annual Field Conference Guidebook, p. 59-62.

Black, B.A., 1984, Oil and Gas Exploration in the Rio Grande Rift, New Mexico: Oil and Gas Fields of the Four Corners Area, p. 799-803.

Black, B.A., 1979, Oil and gas exploration in the Santa Fe-Gallisteo-Hagan area of New Mexico, in New Mexico Geological Society Guidebook, 30th Field conference, Santa Fe Country, p. 275-280.

Broadhead, R.F., 1989, Petroleum production and frontier exploration in New Mexico, in Energy Frontiers in the Rockies, J.C. Lorenz and S.G. Lucas (eds.), , Albuquerque Geological Society, p. 35-46.

Broadhead, R.F., 1996, Oil and gas activities in New Mexico in 1996: New Mexico Geology, v.19, no.4, p. 81-88.

Brooks, B.J., and W.R., Clark, 1978, Wagonmound Dakota-Morrison, in Oil and gas fields of the Four Corners Area, J.E. Fassett, and Thomadis, N.D., eds., Four Corners Geological Society, Colorado.

Butler, W.C., 1988, The rationale for assessment of undiscovered economically recoverable oil and gas in South-Central New Mexico: a geologic overview and play analysis of two favorable areas: U.S. Geological Survey, Open-File Report 88-450-B, 134p.

Chapin, C.E., 1971, The Rio Grande rift, Part.I: modifications and additions: New Mexico Geological Society, Guidebook for the 22nd Field Conference, p. 191-202.

Charpentier, Ronald R., et al., 1996, Tabular Data, Text, and Graphical Images in Support of the 1995 National Assessment of United States Oil and Gas Resources, United States Geological Survey, Digital Data Series DDS-36, CD-ROM.

Clair, J.R., and Brandish, B.B., 1956, Garcia gas field, Las Animas County, Colorado, in Guidebook to the Geology of Raton Basin, Colorado, p. 75-78.

Close, J.C., and Dutcher, R.R., 1993, Predication of permeability trends and origins in coal-bed methane reservoirs of the Raton Basin, New Mexico and Colorado: New Mexico Geological Society, Guidebook to 41st Field Conference, p. 387-395.

Cobban, W.A., and Hook, S.C., 1989, Mid-Cretaceous molluscan biostratigraphy and paleogeography of the southwestern part of the Western Interior, United States, in Westerman, G.E., ed., Jurassic-Cretaceous Biochronology and Paleogeography of North America: Geological Association of Canada Special Paper 27, p. 257-271.

Cordell-Lindrith, Keller, G.R., Hildenbrand, T.G., 1982, Complete Bouguer gravity anomaly map of the Rio Grande Rift, Colorado, New Mexico, and Texas: U.S. Geological Survey Geophysical Investigations Map

Dane, C.H., Landis, E.R., Cobban, W.A., 1971, The Twowells Tongue of the Dakota Sandstone and the Tres Hermanos Sandstone as used by Herrick (1900), Western New Mexico: U.S. Geological Survey Professional Paper 750-B, p. B17-B22.

Dolly, and Meissner, 1977, Oil and gas exploration in North-Central New Mexico, in Exploration frontiers of the Central and Southern Rockies: Rocky Mountain Association of Geologists, p. 23-30.

Eaton, J.G., and Nations, J.G., 1991, Introduction; tectonic setting along the margin of the Cretaceous Western Interior Seaway, southwestern Utah and northern Arizona: Geological Society of America Special Paper 260, p. 1-8.

Gautier, D.L., et al., 1996, 1995 National Assessment of United States Oil and Gas Resources-Results, Methodology, and Supporting Data, United States Geological Survey Digital Data Series DDS-30, Release 2.

Geldon, A.L., 1989, Ground-water hydrology of the central Raton Basin, Colorado and New Mexico: U.S. Geological Survey, Water-supply Paper 2288, 81p. (CBM-NM).

Gilbert, J.L., and Asquith, G.B., 1976, Sedimentology of braided alluvial interval of Dakota Sandstone, northeastern New Mexico: New Mexico Bureau of Mines and Mineral Resources Circular 150, 16p.

Grant, P.R., Jr. and Foster, R.W., 1989, Future petroleum provinces in New Mexico-discovering new reserves: New Mexico Bureau of Mines and Mineral Resources, p. 57-58.

Gries, R.R., 1985, San Juan Sag: Cretaceous rocks in a volcanic-covered basin, South-Central Colorado, The Mountain Geologist, v. 22, no. 4, p. 167-179.

Hook, S.C., 1983, Stratigraphy, paleontology, depositional framework and nomenclature of marine Upper Cretaceous Rocks, Socorro County, New Mexico: New Mexico Geological Society Guidebook, 34th Field Conference, p. 165-172.

Hook, S.C., Cobban, W.A., and Landis, E.R., 1980, Extension of the intertongued Dakota Sandstone-Mancos Shale terminology in the Southern Zuni Basin: New Mexico Geology, v. 2, p. 42-44.

Jacka, A.D., and Brand, J.P., 1972, An analysis of the Dakota Sandstone in the vicinity of Las Vegas, New Mexico, and eastward to the Canadian River Valley: New Mexico Geological Society 23rd Annual Field Conference Guidebook, p. 105-107.

Johnson, R.B., and Wood, G.H., Jr., 1956, Stratigraphy of Upper Cretaceous and Tertiary rocks of Raton Basin, Colorado and New Mexico: AAPG Bulletin, v. 40, p. 707-721.

Johnson, R.E., 1983, Bravo Dome carbon dioxide area, northeast New Mexico, in Oil and gas fields of the Four Corners Area, vol. III, Four Corners Geological Society, p. 745-748.

Keighin, C.W., 1995, Raton Basin-Sierra Grande Uplift Province, in Gautier, D.L., et al., 1996, 1995 National Assessment of United States Oil and Gas Resources-Results, Methodology, and Supporting Data, United States Geological Survey Digital Data Series DDS-30, Release 2.

Kelley, V.C., 1979, Tectonics, middle Rio Grande Rift, New Mexico, in Riecker, R.E., ed., Rio Grande Rift, tectonics and magnetism: American Geophysical Union, Washington, D.C., p. 57-70.

Kelly, V.C., 1978, Geology of the Espanola Basin, New Mexico: New Mexico Bureau of Mines and Mineral Resources, Geologic Map 48.

Kelley, V.C., 1977, Geology of the Albuquerque Basin, New Mexico Bureau of Mines & Mineral Resources, Memoir 33.

Landis, E.R., Dane, C.H., and Cobban, W.A., 1973, Stratigraphic terminology of the Dakota Sandstone and Mancos Shale, West-Central New Mexico: U.S. Geological Survey Bulletin 372, p. J1-J44.

Matuszczak, R.A., 1969, Trinidad Sandstone, interpreted, evaluated, in Raton Basin, Colorado-New Mexico: U.S. Geological Survey Oil and Gas Investigations Preliminary Map 54.

Mellere, D., 1994, Sequential development of an eustarine valley fill: The Twowells Tongue of the Dakota Sandstone, Acoma Basin, New Mexico, Journal of Sedimentary Research, v. B64, no. 4, p. 500-515.

Molenaar, C.M., 1993, Albuquerque-Santa Fe-san Luis Rift Basins Province, in Petroleum exploration plays and resource estimates, 1989, onshore United States; Region 3, Colorado Plateau and Basin and Range., R.B., Powers, ed., U.S. Geological Survey Open-File Report, p. 66-73.

Molenaar, C.M., 1987, Petroleum geology and hydrocarbon plays of the Albuquerque-San Luis Rift Basin, New Mexico and Colorado: U.S. Geological Survey Open-File Report 87-450-S, 26p.

Molenaar, C.M., 1983, Major depositional cycles and regional correlations of Upper Cretaceous rocks, southern Colorado Plateau and adjacent areas, in Reynolds, M.W., and Dolly, E.D., eds., Paleogeography of West-Central United States: SEPM Rocky Mountain Section, p. 201-223.

Molenaar, C.M., 1974, Correlation of the Gallup Sandstone and associated formations, Upper Cretaceous, Eastern San Juan and Acoma Basins, New Mexico, in New Mexico Geological Society Guidebook, 25th Field Conference, Ghost Ranch (Central-Northern NM), p. 251-258.

Northrop, S.A., Sullwold, H.H., Jr., McAlpin, A.J., and Rogers, C.P., Jr., 1946, Geologic maps of a part of the Las Vegas Basin and of the foothills of the Sangre de Cristo Mountains, San Miguel and Mora Counties, New Mexico: U.S. Geological Survey Oil and Gas Investigations Preliminary Map 54.

Nummedal, D., and Riley, G.W., 1991, Origin of Late Turonian and Coniacian unconformities in the San Juan Basin, in Van Wagoner, J.C., Nummedal, D., Jones, C.R., Taylor, D.R., Jennette, D.C., and Riley, G.W., eds., Sequence Stratigraphy Applications to Shelf Sandstone Reservoirs-Outcrop to Subsurface Example: American Association of Petroleum Geologists Field Conference.

Pillmore, C.L., 1969, Geologic map of the Casa Grande quadrangle, Colfax County, New Mexico, and Las Animas County, Colorado: U.S. Geological Survey Geological Quadrangle Map GQ-823.

**Pueblo Tribes of New Mexico**  
REFERENCES (continued)

Pillmore, C.L., 1991, Coal-bed methane in Raton Basin, in coal fields of New Mexico-geology and resources, in U.S. Geological Survey Bulletin 1972, C.L. Molina, ed., p. 15-33, p. 45-68, and p. 69-77.

Rice, D.D., and Finn, T.M., 1995, Coal-bed gas plays in Gautier, D.L., et al., 1996, 1995 National Assessment of United States Oil and Gas Resources-Results, Methodology, and Supporting Data, United States Geological Survey Digital Data Series DDS-30, Release 2.

Roth, G., 1983, Sheep Mountain and Dike Mountain Fields, Huerfano County, Colorado; A source of CO<sub>2</sub> for enhanced oil recovery, in *Oil and Gas Fields of the Four Corners Areas*, J.E. Fassett, ed., Colorado, United States, p. 740-744.

Simms, R.W., 1965, Geology of the Rayando area, Colfax County, New Mexico: Master's thesis, University of New Mexico, Albuquerque, New Mexico, 90p.

Speer, W.R., 1976, Oil and gas exploration in the Raton Basin: New Mexico Geological Society 27th Annual Field Conference Guidebook, p. 217-226.

Stevens, S.H., Lombardi, T.E., Kelso, B.S., and Coates, J.M., 1992, A geologic assessment of natural gas from coal seams in the Raton and Vermejo Formations, Raton Basin. Advanced Resources International, Inc., Report GRI-92/0345, 84p.

Tyler, R., Ambrose, W.A., Scott, A.R., and Kaiser, W.R., 1991, Coal-bed methane potential of the Greater Green River, Piceance, Powder River, and Raton Basins: Bureau of Economic Geology, University of Texas at Austin, Report GRI-91/00315, 249 p.

Van Wagoner, J.C., Mitchum, R.M., Campion, K.M., and Ramani, V.D., 1990, Siliclastic sequence stratigraphy in well logs, cores, and outcrops: techniques for high resolution correlation of time and facies, American Association of Petroleum Geologists Methods in Exploration Series no. 7, 55p.

Van Wagoner, J.C., Mitchum, R.M., Posamentier, H.W., and Vail, P.R., 1988, An overview of fundamentals of sequence stratigraphy and key definitions of sequence stratigraphy, in Wilgus, C.K., Hastings, B.S., Kendall, C.G.St.C., Posamentier, H.W., Ross, C.A., and Van Wagoner, J.C. (eds.), *Sea Level Changes: an Integrated Approach*: SEPM Special Publications 42, p. 38-47.

Villien, A., and Kligfield, R.M., 1986, Thrusting and synorogenic sedimentation in central Utah, in Peterson, J.A., ed., *Paleotectonics and Sedimentation in the Rocky Mountain Region, United States*: American Association of Petroleum Geologists Memoir 41, p. 281-308.

Woodward, L.A., and Snyder, D.O., 1976, Tectonic map of the southern Raton Basin, New Mexico: New Mexico Geological Society 27th Annual Field Conference Guidebook, Map.

Woodward, L.A., 1974, Tectonics of Central-Northern New Mexico, New Mexico Geological Society Guidebook, 25th Field Conference, Ghost Ranch (Central-Northern NM), p. 123-129.

Woodward, L.A., 1984, Geology and Hydrocarbon Potential of the Raton Basin, New Mexico: Oil and Gas Fields of the Four Corners Area, p. 789-798.